

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all of the information needed to use CEFTRIAXONE FOR INJECTION safely and effectively. See full prescribing information for CEFTRIAXONE FOR INJECTION.

CEFTRIAXONE FOR INJECTION, USP

Ceftriaxone for Injection, for intravenous use

PHARMACY BULK PACKAGE – NOT FOR DIRECT INFUSION
--

Initial U.S. Approval: 1984

To reduce the development of drug-resistant bacteria and maintain the effectiveness of ceftriaxone and other antibacterial drugs, ceftriaxone should be used only to treat or prevent infections that are proven or strongly suspected to be caused by bacteria.

INDICATIONS AND USAGE

Ceftriaxone for Injection, USP is a cephalosporin antibacterial indicated for the treatment of the following infections caused by susceptible isolates of the designated bacteria: Lower Respiratory Tract Infections (1.1); Skin and Skin Structure Infections (1.2); Complicated and Uncomplicated Urinary Tract Infections (1.3); Pelvic Inflammatory Disease (1.4); Bacterial Septicemia (1.5); Bone and Joint Infections (1.6); Intra-abdominal Infections (1.7); Meningitis (1.8); and Surgical Prophylaxis (1.9).

DOSAGE AND ADMINISTRATION

For intravenous use only over approximately 30 minutes. (2)
THIS IS A PHARMACY BULK PACKAGE – NOT FOR DIRECT INJECTION.

Ceftriaxone for Injection USP, Pharmacy Bulk Package bag SmartPak® should be used only in patients who require a 1 gram dose and not any fraction thereof. Ceftriaxone for Injection USP, Pharmacy Bulk Package bag SmartPak® should not be used in patients who require less than the 1 gram dose of ceftriaxone. (2.1).

Recommended Dosing Schedule for Ceftriaxone for Injection, USP			
Site and Type of Infection	Dose	Frequency	Total Daily Dose
Usual Adult Dose	1 to 2 grams	Once a day or in equally divided doses every 12 hours	Should not exceed 4 grams*
Surgical prophylaxis	1 gram I.V. once	½ to 2 hours before surgery	
Meningitis	100 mg per kg	Once a day or in equally divided doses every 12 hours	Should not exceed 4 grams*
Skin and Skin Structure Infections	50 mg per kg to 75 mg per kg	Once a day or in equally divided doses every 12 hours	Should not exceed 2 grams
Serious Infections other than Meningitis	50 mg per kg to 75 mg per kg	Every 12 hours	Should not exceed 2 grams

*Patients with hepatic impairment and significant renal impairment should not receive more than 2 grams per day of ceftriaxone.

Ceftriaxone for Injection USP, Pharmacy Bulk Package bag SmartPak® should be used only in patients who require a 1 gram dose and not any fraction thereof. Ceftriaxone for Injection USP, Pharmacy Bulk Package bag SmartPak® should not be used in patients who require less than the 1 gram dose of ceftriaxone. (2.1)

DOSAGE FORMS AND STRENGTHS

Pharmacy Bulk Package bags, 100 grams (3)

THIS IS A PHARMACY BULK PACKAGE – NOT FOR DIRECT INJECTION

- Anaphylaxis to ceftriaxone or other cephalosporin class antibacterials, penicillins or other beta-lactam antibacterials (4.1).
- WARNINGS AND PRECAUTIONS**
- Hypersensitivity reactions: Include anaphylaxis and serious skin reactions. Cross hypersensitivity may occur in up to 10% of patients with a history of penicillin allergy. If an allergic reaction occurs, discontinue the drug (5.1).
- Interaction with Calcium-containing products: Precipitation can occur. Do not administer simultaneously with calcium-containing I.V. solutions (5.2).
- Neurological adverse reactions: Serious neurological adverse reactions have been reported during postmarketing surveillance. If serious adverse reactions occur, discontinue the drug and institute appropriate supportive measures. (5.3)
- Clostridium difficile*-associated diarrhea: May range from mild diarrhea to fatal colitis. Evaluate if diarrhea occurs (5.4).
- Hemolytic Anemia: Severe cases of hemolytic anemia, including fatalities in adults and children, have been reported. If anemia is diagnosed, discontinue the drug until the etiology is determined. (5.5)


ADVERSE REACTIONS

The most common adverse reactions occurring in >2% of patients receiving ceftriaxone include diarrhea, eosinophilia, thrombocytosis, leukopenia, and elevations of SGOT and SGPT.
To report SUSPECTED ADVERSE REACTIONS, contact Samson Medical Technologies, L.L.C. at 1-877-418-3600 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch

1

INSTRUCTION FOR RECONSTITUTION OF THE PHARMACY BULK PACKAGE BAG SmartPak®

The entire contents of the bag and the preparation process (reconstitution and dilution) should be completed within 4 hours of initial entry.



- Document the date and time reconstitution starts in the designated place on the container label. The entire contents of the bag must be used within **4 hours** from the time of initial entry.
- Remove the cap from the reconstitution (smaller) port and discard it.
- Reconstitute the powder through the reconstitution (smaller) port, using Sterile Water for Injection according to the Table 2 below.

Table 2: Reconstitution Table		
SmartPak® Bag Size	Amount of Sterile Water*	Approximate Concentration
100 grams	950 mL	100 mg/mL (1 g/10 mL)

***Do not use diluents containing calcium, such as Ringer's solution or Hartmann's solution, to reconstitute or further dilute reconstituted ceftriaxone because a precipitate can form. STERILE WATER FOR INJECTION IS THE ONLY RECOMMENDED DILUENT FOR RECONSTITUTION.** [see *Warnings and Precautions* (5.2)].

- After reconstitution is complete, remove the transfer needle from the reconstitution port
- Place the bag on a flat surface of a laminar flow hood and mix for at least 15 minutes by rocking gently from side to side. **CAUTION: To avoid possible leakage caused by the heavy weight of the added water, do not shake vigorously or pull strongly on the bag.**
- When foam dissipates, visually inspect the bag to verify the solution is clear, colorless to pale yellow and free of particulate matter. **DO NOT USE THE INNER BAG IF PARTICULATE OR FOREIGN MATTER IS PRESENT.**
- Remove the cap from the transfer (larger) port and discard it. Attach sterile tubing and filling adapter unit to the transfer port.
- Reconstituted solution can now be transferred using the transfer port and the filling adapter.

It should be noted that the spike placed into the transfer port of the Pharmacy Bulk Package SmartPak® is NEVER removed during this procedure and that the reconstitution port is self-sealing.

5

DRUG INTERACTIONS

- Vancomycin, amasacrine, aminoglycosides and fluconazole are physically incompatible. (7.1)
- Calcium-containing products: precipitation can occur (7.2).

USE IN SPECIFIC POPULATIONS

- Hepatic impairment:** Patients with both hepatic and renal impairment should not receive more than 2 grams of ceftriaxone per day (5.7).
- Renal Impairment:** This formulation of Ceftriaxone for Injection USP – Pharmacy Bulk Package bag, SmartPak® should not be used in renally impaired patients who require less than the 1 gram dose of ceftriaxone. Patients with both hepatic and renal impairment should not receive more than 2 grams of ceftriaxone per day. (5.7)
- Pediatric Patients:** This formulation of Ceftriaxone for Injection USP – Pharmacy Bulk Package bag, SmartPak® should not be used in pediatric patients who require less than the 1 gram adult dose of ceftriaxone (2.2, 8.4).

See 17 for PATIENT COUNSELING INFORMATION

Revised 04/2023

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

- Lower Respiratory Tract Infections
- Skin and Skin Structure Infections
- Complicated and Uncomplicated Urinary Tract Infections
- Pelvic Inflammatory Disease
- Bacterial Septicemia
- Bone and Joint Infections
- Intra-Abdominal Infections
- Meningitis
- Surgical Prophylaxis

2 DOSAGE AND ADMINISTRATION

- Adult Population
- Pediatric Population
- Preparation for Use of Ceftriaxone for Injection, USP, Pharmacy Bulk Package bags, SmartPak®

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

- Anaphylaxis to Ceftriaxone or the Cephalosporin Class of Antibacterials, Penicillins or Other Beta-lactam Antibacterials

5 WARNINGS AND PRECAUTIONS

- Hypersensitivity Reactions to Ceftriaxone, Cephalosporins, Penicillins or Other Drugs
- Interaction with Calcium-Containing Products
- Neurological Adverse Reactions
- Bone and joint infections caused by *Staphylococcus aureus*, *Streptococcus pneumoniae*, *Escherichia coli*, *Proteus mirabilis*, *Klebsiella pneumoniae* or *Enterobacter* species.
- Clostridium difficile*-associated Diarrhea
- Hemolytic Anemia
- Gallbladder Sonogram Abnormalities
- Patients with Hepatic and Renal Impairment
- Pancreatitis
- Development of Drug-resistant Bacteria
- Alterations in Prothrombin Time

6 ADVERSE REACTIONS

- Clinical Trials Experience
- Post-Marketing Experience
- Cephalosporin–class Adverse Reactions

7 DRUG INTERACTIONS

- Vancomycin, Amasacrine, Aminoglycosides, and Fluconazole
- Calcium-containing Products

8 USE IN SPECIFIC POPULATIONS

- Pregnancy
- Nursing Mothers
- Pediatric Use
- Geriatric Use

10 OVERDOSAGE

11 DESCRIPTION

- CLINICAL PHARMACOLOGY
 - Mechanism of Action
 - Pharmacokinetics
 - Microbiology

13 NON-CLINICAL TOXICOLOGY

- Carcinogenesis, Mutagenesis, Impairment of Fertility
- Animal Toxicology and/or Animal Pharmacology

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

*Sections or subsections omitted from the full prescribing information are not listed.

2

Dilution

- Hang the bag from two eyelets.
- Following reconstitution, transfer 10 mL of the reconstituted solution into transfusion bags, each containing 50 mL of one of the compatible solutions below.

Compatible solutions for dilution are the following:

- Sodium Chloride Injection, USP
- 5% Dextrose Injection, USP

- Dilution should be completed within the 4 hour preparation process.
- When diluted according to the instructions above, ceftriaxone is stable for 48 hours at room temperature or for 10 days if stored under refrigeration (5°C or 41°F).

Important Administration Instructions

- Do not use diluents containing calcium, such as Ringer's solution or Hartmann's solution, to reconstitute Ceftriaxone for Injection SmartPak® Pharmacy Bulk Package or to further dilute a reconstituted solution for intravenous administration. Particulate formation can result.** Precipitation of ceftriaxone-calcium can also occur when ceftriaxone is mixed with calcium-containing solutions in the same intravenous administration line. Ceftriaxone for Injection, USP must not be administered simultaneously with calcium-containing intravenous solutions, including continuous calcium-containing infusions, such as parenteral nutrition via a Y-site. However, in patients other than neonates, Ceftriaxone for Injection, USP and calcium-containing solutions may be administered sequentially of one another if the infusion lines are thoroughly flushed between infusions with a compatible fluid [see *Warnings and Precautions* (5.2)].
- Prior to administration, parental products should be inspected visually for particulate matter and discoloration whenever solution and container permit.
- Administer diluted ceftriaxone solutions intravenously over approximately 30 minutes.
- After the indicated stability times, unused portions of solutions should be discarded.
- Vancomycin, amasacrine, aminoglycosides and fluconazole are physically incompatible with ceftriaxone in admixtures. When any of these drugs are to be administered concomitantly with ceftriaxone by intermittent intravenous infusion, it is recommended that they be given sequentially with thorough flushing of the intravenous lines with 0.9% Sodium Chloride Injection or 5% Dextrose in Water (D5W) between the administrations.
- Ceftriaxone for Injection should not be physically mixed with or piggybacked into solutions containing other antimicrobial drugs due to possible incompatibility [see *Drug Interactions* (7.1)].
- There have been no reports of an interaction between ceftriaxone and oral calcium-containing products or interaction between intramuscular ceftriaxone and calcium-containing products (intravenous or oral).

DOSAGE FORMS AND STRENGTHS

- 100 grams Ceftriaxone for Injection, USP, Pharmacy Bulk Package bag, SmartPak®

THIS IS A PHARMACY BULK PACKAGE – NOT FOR DIRECT INJECTION

CONTRAINDICATIONS

4.1 Anaphylaxis to Ceftriaxone or the Cephalosporin Class of Antibacterials, Penicillins or Other Beta-Lactam Antibacterials
Ceftriaxone for Injection, USP is contraindicated in patients who have a history of anaphylaxis to ceftriaxone or the cephalosporin class of antibacterials, penicillins, or other beta-lactam antibacterials [see *Warnings and Precautions* (5.1)].

WARNINGS AND PRECAUTIONS

5.1 Hypersensitivity Reactions to Cephalosporins, Penicillins, or Other Drugs

Serious, occasionally fatal, hypersensitivity (anaphylactic) reactions have been reported with ceftriaxone. Before therapy with Ceftriaxone for Injection, USP is instituted, careful inquiry should be made to determine whether the patient has had previous immediate hypersensitivity reactions to ceftriaxone, cephalosporins, penicillins, or other drugs. Exercise caution if this product is to be given to penicillin-sensitive patients because cross-hypersensitivity among beta-lactam antibacterials has been clearly documented and may occur in up to 10% of patients with a history of penicillin allergy. If an allergic reaction to Ceftriaxone for Injection, USP occurs, discontinue the drug. Serious acute hypersensitivity reactions may require treatment with epinephrine and other emergency measures including oxygen, corticosteroids, intravenous fluids, intravenous antihistamines, pressor amines, and airway management, as clinically indicated.

FULL PRESCRIBING INFORMATION

INDICATIONS AND USAGE

To reduce the development of drug-resistant bacteria and maintain the effectiveness of Ceftriaxone for Injection and other antibacterial drugs, Ceftriaxone for Injection should be used only to treat or prevent infections that are proven or strongly suspected to be caused by susceptible bacteria. When culture and susceptibility information are available, they should be considered in selecting or modifying antibacterial therapy. In the absence of such data, local epidemiology and susceptibility patterns may contribute to the empiric selection of therapy.

Ceftriaxone for Injection is indicated for the treatment of the following infections when caused by susceptible bacteria:

1.1 Lower Respiratory Tract Infections

Lower respiratory tract infections caused by *Streptococcus pneumoniae*, *Staphylococcus aureus*, *Haemophilus influenzae*, *Haemophilus parainfluenzae*, *Klebsiella pneumoniae*, *Escherichia coli*, *Enterobacter aerogenes*, *Proteus mirabilis* or *Serratia marcescens*.

1.2 Skin and Skin Structure Infections

Skin and skin structure infections caused by *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Streptococcus pyogenes*, Viridans group streptococci, *Escherichia coli*, *Enterobacter cloacae*, *Klebsiella oxytoca*, *Klebsiella pneumoniae*, *Proteus mirabilis*, *Morganella morganii*,* *Pseudomonas aeruginosa*, *Serratia marcescens*, *Acinetobacter calcoaceticus*, *Bacteroides fragilis** or *Peptostreptococcus* species.

*The efficacy for these organisms in this organ system was studied in fewer than ten infections.

1.3 Complicated and Uncomplicated Urinary Tract Infections

Complicated and uncomplicated urinary tract infections caused by *Escherichia coli*, *Proteus mirabilis*, *Proteus vulgaris*, *Morganella morganii* or *Klebsiella pneumoniae*.

1.4 Pelvic Inflammatory Disease

Pelvic inflammatory disease caused by *Neisseria gonorrhoeae*. Ceftriaxone sodium, like other cephalosporins, has no activity against *Chlamydia trachomatis*. Therefore, when cephalosporins are used in the treatment of patients with pelvic inflammatory disease and *Chlamydia trachomatis* is one of the suspected pathogens, appropriate antichlamydial coverage should be added.

1.5 Bacterial Septicemia

Bacterial septicemia caused by *Staphylococcus aureus*, *Streptococcus pneumoniae*, *Escherichia coli*, *Haemophilus influenzae* or *Klebsiella pneumoniae*.

1.6 Bone and Joint Infections

Bone and joint infections caused by *Staphylococcus aureus*, *Streptococcus pneumoniae*, *Escherichia coli*, *Proteus mirabilis*, *Klebsiella pneumoniae* or *Enterobacter* species.

1.7 Intra-abdominal Infections

Intra-abdominal infections caused by *Escherichia coli*, *Klebsiella pneumoniae*, *Bacteroides fragilis*, *Clostridium* species or *Peptostreptococcus* species.

1.8 Meningitis

Meningitis caused by *Haemophilus influenzae*, *Neisseria meningitidis* or *Streptococcus pneumoniae*. Ceftriaxone sodium has also been used successfully in a limited number of cases of meningitis and shunt infection caused by *Staphylococcus epidermidis* and *Escherichia coli*; however, the efficacy for these organisms in this organ system were studied in fewer than ten infections.

1.9 Surgical Prophylaxis

The preoperative administration of a single 1 gram dose of ceftriaxone may reduce the incidence of postoperative infections in patients undergoing surgical procedures classified as contaminated or potentially contaminated (e.g., vaginal or abdominal hysterectomy or cholecystectomy for chronic calculous cholecystitis in high-risk patients, such as those over 70 years of age, with acute cholecystitis not requiring therapeutic antimicrobials, obstructive jaundice or common duct bile stones) and in surgical patients for whom infection at the operative site would present serious risk (e.g., during coronary artery bypass surgery). Although ceftriaxone sodium has been shown to have been as effective as cefazolin in the prevention of infection following coronary artery bypass surgery, no placebo-controlled trials have been conducted to evaluate any cephalosporin antibacterial in the prevention of infection following coronary artery bypass surgery.

DOSAGE AND ADMINISTRATION

Ceftriaxone for Injection USP, Pharmacy Bulk Package bag SmartPak® should be used only in patients who require a 1 gram dose and not any fraction thereof.

Ceftriaxone for Injection USP, Pharmacy Bulk Package bag SmartPak® should not be used in patients who require less than the 1 gram dose of Ceftriaxone.

THIS PHARMACY BULK PACKAGE REQUIRES RECONSTITUTION WITH STERILE WATER FOR INJECTION, USP TO A CONCENTRATION OF 100 mg per mL AND FURTHER DILUTION IN 50 mL OF A COMPATIBLE SOLUTION.

THIS IS A PHARMACY BULK PACKAGE – NOT FOR DIRECT INJECTION

USE THIS FORMULATION OF CEFTRIAXONE ONLY IN PATIENTS WHO REQUIRE A 1 GRAM DOSE.

3

5.2

Interaction with Calcium-Containing Products

Precipitation of ceftriaxone-calcium can occur when Ceftriaxone for Injection, USP is mixed with calcium-containing solutions in the same intravenous administration line. Ceftriaxone for Injection must not be administered simultaneously with calcium-containing intravenous solutions, including continuous calcium-containing infusions, such as parenteral nutrition via a Y-site. However, in patients other than neonates, Ceftriaxone for Injection, USP and calcium-containing solutions may be administered sequentially of one another if the infusion lines are thoroughly flushed between infusions with 0.9% Sodium Chloride Injection or D5W. *In vitro* studies using adult and neonatal plasma from umbilical cord blood demonstrated that neonates have an increased risk of precipitation of ceftriaxone-calcium. [see *Drug Interactions* (7.2)]

5.3 Neurological Adverse Reactions

Serious neurological adverse reactions have been reported during postmarketing surveillance with ceftriaxone use. These reactions include encephalopathy (disturbances of consciousness including somnolence, lethargy, and confusion), seizures, myoclonus, and non-convulsive status epilepticus [see *Adverse Reactions* (6.2)]. Some cases occurred in patients with severe renal impairment who did not receive appropriate dosage adjustment. However, in other cases, neurological adverse reactions occurred in patients receiving an appropriate dosage adjustment. The neurological adverse reactions were reversible and resolved after discontinuation. If neurological adverse reactions associated with Ceftriaxone for Injection therapy occur, discontinue Ceftriaxone for Injection and institute appropriate supportive measures. Make appropriate dosage adjustments in patients with severe renal impairment. [see *Dosage and Administration* (2.1)].

5.4 *Clostridium difficile*-associated Diarrhea

Clostridium difficile-associated diarrhea (CDAD) has been reported with use of nearly all antibacterial agents, including ceftriaxone, and may range in severity from mild diarrhea to fatal colitis. Treatment with antibacterial agents alters the normal flora of the colon leading to overgrowth of *C. difficile*.

C. difficile produces toxins A and B, which contribute to the development of CDAD. Hypertoxin-producing strains of *C. difficile* cause increased morbidity and mortality, as these infections can be refractory to antimicrobial therapy and may require colectomy. CDAD must be considered in all patients who present with diarrhea following antibacterial use. Careful medical history is necessary, since CDAD has been reported to occur over two months after the administration of antibacterial agents.

If CDAD is suspected or confirmed, ongoing antibacterial use not directed against *C. difficile* may need to be discontinued. Appropriate fluid and electrolyte management, protein supplementation, antibacterial treatment of *C. difficile*, and surgical evaluation should be instituted as clinically indicated.

5.5 Hemolytic Anemia

An immune mediated hemolytic anemia has been observed in patients receiving cephalosporin class antibacterials including ceftriaxone. Severe cases of hemolytic anemia, including fatalities, have been reported during treatment in both adults and children. If a patient develops anemia while on Ceftriaxone for Injection, USP, the diagnosis of a cephalosporin-associated anemia should be considered and Ceftriaxone for Injection, USP stopped until the etiology is determined.

5.6 Gallbladder Sonogram Abnormalities

There have been reports of sonographic abnormalities in the gallbladder of patients treated with ceftriaxone sodium; some of these patients also had symptoms of gallbladder disease. These abnormalities appear on sonography as an echo without acoustical shadowing suggesting sludge or as an echo with acoustical shadowing, which may be misinterpreted as gallstones. The chemical nature of the sonographically detected material has been determined to be predominantly a ceftriaxone-calcium salt. The condition appears to be transient and reversible upon discontinuation of ceftriaxone sodium and institution of conservative management. Therefore, ceftriaxone should be discontinued in patients who develop signs and symptoms suggestive of gallbladder disease and/or the sonographic findings described above.

5.7 Patients with Hepatic and Renal Impairment

Hepatic Impairment: In patients with both hepatic and significant renal disease, Ceftriaxone for Injection dosage should not exceed 2 grams daily.

Renal Impairment: This formulation of Ceftriaxone for Injection USP – Pharmacy Bulk Package bags SmartPak® should not be used in renally impaired patients who require less than the 1 gram dose of ceftriaxone. In patients with both hepatic impairment and significant renal disease, Ceftriaxone for Injection dosage should not exceed 2 grams per day.

2.1 Adult Population

Ceftriaxone for Injection USP, Pharmacy Bulk Package bag SmartPak® should be used only in patients who require a 1 gram dose and not any fraction thereof.

Ceftriaxone for Injection USP Pharmacy Bulk Package bag SmartPak® should not be used in patients who require less than the 1 gram dose of Ceftriaxone.

Ceftriaxone for Injection should be reconstituted with Sterile Water for Injection, USP to a concentration of 100 mg per mL and further diluted in 50 mL of a compatible solution.

The recommended adult dosages are outlined in Table 1. Ceftriaxone for Injection USP, Pharmacy Bulk Package SmartPak® should be used only in patients who require a 1 gram dose and not any fraction thereof.

Ceftriaxone should be administered intravenously (I.V.) over approximately 30 minutes.

The usual duration of therapy is 4 to 14 days; in complicated infections, longer therapy may be required. When treating infections caused by *Streptococcus pyogenes*, therapy should be continued for at least 10 days.

Table 1: Recommended Dosing Schedule for Ceftriaxone for Injection, USP			
Site and Type of Infection	Dose	Frequency	Total Daily Dose
Usual Adult Dose	1 gram to 2 grams	Once a day or in equally divided doses every 12 hours	Should not exceed 4 grams*
Surgical Prophylaxis	1 gram I.V. once	½ to 2 hours before surgery	
Skin and Skin Structure Infections	50 to 75 mg per kg	Once a day or in equally divided doses every 12 hours	Should not exceed 2 grams
Meningitis	100 mg per kg	Once a day or in equally divided doses every 12 hours	Should not exceed 4 grams*
Serious Infections other than Meningitis	50 to 75 mg per kg	Every 12 hours	Should not exceed 2 grams

*Patients with hepatic impairment and significant renal impairment should not receive more than 2 grams per day of ceftriaxone.
****This formulation of Ceftriaxone for Injection USP, Pharmacy Bulk Package bag SmartPak® should not be used in patients who require less than the 1 gram dose of ceftriaxone.**

2.2 Pediatric Patients

To prevent unintentional overdose, this product should not be used in pediatric patients who require less than a 1 gram dose of ceftriaxone. [see *Use in Specific Populations* (8.4)]

2.3 Preparation for Use of Ceftriaxone for Injection, USP, bag SmartPak® Pharmacy Bulk Package

Ceftriaxone for Injection, USP Pharmacy Bulk Package bag SmartPak® should be used only in patients who require a 1 gram dose and not any fraction thereof.

Ceftriaxone for Injection USP, Pharmacy Bulk Package SmartPak® bag should not be used in patients who require less than the 1 gram dose of ceftriaxone.

Directions for Proper Use of a Pharmacy Bulk Package

- NOT FOR DIRECT INFUSION.** The Pharmacy Bulk Package is for use in the hospital pharmacy admixture service only in a suitable work area, such as a laminar flow hood. Using aseptic technique, the container closure may be penetrated only one time after reconstitution using a suitable sterile dispensing set or transfer device that allows measured dispensing of the contents. Use of a syringe and needle is not recommended as it may cause leakage. The withdrawal of container contents should be accomplished without delay. However, should this not be possible, a maximum time of 4 **HOURS** from initial reconstitution port closure entry is permitted to complete fluid transfer operations. This time limit should begin with the introduction of the solvent or diluent into the Pharmacy Bulk Package. Discard any unused portion after 4 **HOURS**. This pharmacy bulk package is not intended to be dispensed as a unit
- PRIOR TO RECONSTITUTION: DO NOT USE THE INNER BAG IF PARTICULATE OR FOREIGN MATTER IS PRESENT, IF THE DRY POWDER IS DARK YELLOW OR BROWN, IF THE SEALS ARE NOT INTACT, OR IF THERE IS ANY OTHER DAMAGE TO THE BAG. IN SUCH CASES, DISCARD THE BAG IMMEDIATELY.**
- After initial reconstitution port entry, use entire contents of the Pharmacy Bulk Package promptly. Any unused portion must be discarded after 4 **HOURS**.
- Gather the following items prior to the reconstitution of the product: Appropriate number of bags of Sterile Water for Injection and, depending upon the method of filling, appropriate sterile tubing and adapters.
- Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration. Use only if solution is clear.
- Do not use diluents containing calcium, such as Ringer's solution or Hartmann's solution, to reconstitute Ceftriaxone for Injection SmartPak® Pharmacy Bulk Package or to further dilute a reconstituted solution for intravenous administration.** Particulate formation can result.

Anaphylaxis (including anaphylactic shock, transient leukopenia, neutropenia, agranulocytosis and thrombocytopenia) has been reported.

6.3 Cephalosporin Class Adverse Reactions

In addition to the adverse reactions listed above that have been observed in patients treated with ceftriaxone, the following adverse reactions and altered laboratory tests have been reported for cephalosporin-class antibacterials:

- Adverse Reactions: Allergic reactions, drug fever, serum sickness-like reaction, renal dysfunction, toxic nephropathy, reversible hyperactivity, hypertonia, hepatic dysfunction including cholestasis, aplastic anemia, hemorrhage, and superinfection.
- Altered Laboratory Tests: Positive direct Coombs' test, false-positive test for urinary glucose, and elevated LDH.

7 DRUG INTERACTIONS

7.1 Vancomycin, Amsacrine, Aminoglycosides and Fluconazole

Vancomycin, amsacrine, aminoglycosides, and fluconazole are physically incompatible with ceftriaxone in admixtures [see *Dosage and Administration* (2.3)].

7.2 Calcium-containing Products

Precipitation of ceftriaxone-calcium can occur when Ceftriaxone for Injection is mixed with calcium-containing solutions in the same intravenous administration line. Ceftriaxone for Injection, USP must not be administered simultaneously with calcium-containing intravenous solutions. Ceftriaxone for Injection and calcium-containing solutions can be administered sequentially [see *Warnings and Precautions* (5.2)].

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Pregnancy Category B

Reproductive studies have been performed in mice, rats, and primates at intravenous doses of 625, 586, and 84 mg/kg/day, respectively without evidence of embryotoxicity, fetotoxicity, or teratogenicity. These doses are approximately 1.5, 2.8, and 0.8 times the recommended clinical dose of 2 grams/day based on body surface area comparisons.

There are, however, no adequate and well-controlled studies in pregnant women. Because animal reproductive studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed.

Ceftriaxone was tested in a Segment III (pre-postnatal) study in rats at intravenous doses of up to 586 mg/kg/day approximately 2.8 times (mg/m² comparison) the recommended daily dose of 2 grams/day. No adverse effects were noted on various reproductive parameters during gestation and lactation, including postnatal growth, functional behavior, and reproductive ability of the offspring.

8.3 Nursing Mothers

Ceftriaxone is excreted in human breast milk. Caution should be exercised when Ceftriaxone for Injection is administered to a nursing woman.

8.4 Pediatric Use

Ceftriaxone for Injection USP, Pharmacy Bulk Package bag, SmartPak® should not be used in pediatric patients who require less than the 1 gram dose of ceftriaxone. To prevent unintentional overdose, this product should not be used in pediatric patients who require less than the 1 gram adult dose of ceftriaxone.

8.5 Geriatric Use

Of the total number of subjects in clinical studies of ceftriaxone sodium, 32% were 60 and over. No overall differences in safety or effectiveness were observed between these subjects and younger subjects, and other reported clinical experience has not identified differences in responses between the elderly and younger patients, but greater sensitivity of some older individuals cannot be ruled out.

Ceftriaxone is known to be substantially excreted by the kidney, and the risk of adverse reactions to this drug may be greater in patients with impaired renal function. Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection, and it may be useful to monitor renal function.

The pharmacokinetics of ceftriaxone were only minimally altered in geriatric patients compared to healthy adult subjects, and dosage adjustments are not necessary for geriatric patients with ceftriaxone dosages up to 2 grams per day [see *Clinical Pharmacology* (12)].

10 OVERDOSAGE

Ceftriaxone overdosage has been reported in patients with severe renal impairment. Reactions have included neurological outcomes, including encephalopathy, seizures, myoclonus, and non-convulsive status epilepticus. In the event of overdosage, discontinue Ceftriaxone for Injection therapy and provide general supportive treatment [see *Dosage and Administration* (2.1) and *Warnings and Precautions* (5.3)].

In the case of overdosage, drug concentration would not be reduced by hemodialysis or peritoneal dialysis. There is no specific antidote. Treatment of overdosage should be symptomatic.

9
<ul style="list-style-type: none">Gram-positive bacteria <ul style="list-style-type: none"><i>Staphylococcus aureus</i> <i>Staphylococcus epidermidis</i> <i>Streptococcus pneumoniae</i> <i>Streptococcus pyogenes</i> <i>Viridans group streptococci</i> Anaerobic bacteria <ul style="list-style-type: none"><i>Bacteroides fragilis</i> <i>Clostridium</i> species <i>Peptostreptococcus</i> species

The following *in vitro* data are available, **but their clinical significance is unknown**. At least 90 percent of the following microorganisms exhibit an *in vitro* minimum inhibitory concentration (MIC) less than or equal to the susceptible breakpoint for ceftriaxone. However, the efficacy of ceftriaxone in treating clinical infections due to these microorganisms **has not been** established in adequate and well-controlled clinical trials.

- Gram-negative bacteria**
 - Citrobacter diversus*
 - Citrobacter freundii*
 - Providencia* species (including *Providencia rettgeri*)
 - Salmonella* species (including *Salmonella typhi*)
 - Shigella* species
- Gram-positive bacteria**
 - Streptococcus agalactiae*
- Anaerobic bacteria**
 - Porphyromonas (Bacteroides) melaninogenicus*
 - Prevotella (Bacteroides) bivia*

Susceptibility Testing

For specific information regarding susceptibility test interpretive criteria and associated test methods and quality control standards recognized by FDA for this drug, please see: <https://www.fda.gov/STIC>.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis

Considering the maximum duration of treatment and the class of the compound, carcinogenicity studies with ceftriaxone in animals have not been performed. The maximum duration of animal toxicity studies was 6 months.

Mutagenesis

Genetic toxicology tests included the Ames test, a micronucleus test and a test for chromosomal aberrations in human lymphocytes cultured *in vitro* with ceftriaxone. Ceftriaxone showed no potential for genotoxic activity in these studies.

Impairment of Fertility

Ceftriaxone produced no impairment of fertility when given intravenously to rats at daily doses up to 586 mg/kg/day, approximately 2.8 times (mg/m² comparison) the recommended clinical dose of 2 grams/day.


11 DESCRIPTION

Ceftriaxone for Injection USP, Pharmacy Bulk Package bag SmartPak® should be used only in patients who require a 1 gram dose and not any fraction thereof.

Ceftriaxone for Injection USP, Pharmacy Bulk Package bag SmartPak® should not be used in patients who require less than the 1 gram dose of ceftriaxone.

Ceftriaxone for Injection, USP, is a sterile, semisynthetic, broad-spectrum cephalosporin antibacterial for intravenous administration. Ceftriaxone sodium is (6*R*,7*R*)-7-[2-(2-Amino-4-thiazolyl)glyoxylamido]-8-oxo-3[[[(1,2,5,6-tetrahydro-2-methyl-5,6-dioxo-as-triazin-3-yl)thio]methyl]-5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylic acid, 7⁻ -(Z)-(O-methylxime), disodium salt, sesquaterhydrate.

The chemical formula of ceftriaxone sodium is C₁₈H₁₆N₆Na₂O₇S₃•3.5H₂O. It has a calculated molecular weight of 661.60 and the following structural formula:



Ceftriaxone sodium is a white to yellowish-orange crystalline powder, which is readily soluble in water, sparingly soluble in methanol and very slightly soluble in ethanol. The pH of a 1% aqueous solution is approximately 6.7. The color of ceftriaxone sodium solutions ranges from light yellow to amber, depending on the length of storage and concentration. Each Pharmacy Bulk Package contains approximately 83 mg (3.6 mEq) of sodium per 1 gram of ceftriaxone activity. Ceftriaxone for Injection, USP is supplied in 100 grams SmartPak® Pharmacy Bulk Packages bags equivalent. Each SmartPak® Pharmacy Bulk Package bag contains ceftriaxone sodium, USP equivalent to 100 grams of ceftriaxone.

BEFORE ADMINISTRATION, THIS PHARMACY BULK PACKAGE REQUIRES RECONSTITUTION USING STERILE WATER FOR INJECTION, USP TO A CONCENTRATION OF 100 mg per mL AND FURTHER DILUTION IN 50 mL OF A COMPATIBLE SOLUTION AND INFUSED INTRAVENOUSLY OVER 30 MINUTES.

THIS PRODUCT IS NOT INTENDED TO BE USED IN PEDIATRIC PATIENTS AND RENALLY IMPAIRED PATIENTS WHO REQUIRE LESS THAN A 1 GRAM DOSE.

A Pharmacy Bulk Package is a container of a sterile preparation for parenteral use that contains many single doses. The contents are intended for use in a pharmacy admixture service and are restricted to the preparation of admixtures for intravenous infusion.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Ceftriaxone is an antibacterial drug. [see *Microbiology* (12.4)]

12.3 Pharmacokinetics

Average plasma concentrations of ceftriaxone following a single 30-minute intravenous (I.V.) infusion of a 0.5, 1 or 2 g dose in healthy subjects are presented in Table 3. Multiple intravenous doses ranging from 0.5 to 2 g at 12- to 24-hour intervals resulted in 15% to 36% accumulation of ceftriaxone above single dose values.

Dose/Route	Average Plasma Concentrations (mcg/mL)								
	0.5 hr	1 hr	2 hr	4 hr	6 hr	8 hr	12 hr	16 hr	24 hr
0.5 gram I.V.*	82	59	48	37	29	23	15	10	5
1 gram I.V.*	151	111	88	67	53	43	28	18	9
2 grams I.V.*	257	192	154	117	89	74	46	31	15

*I.V. doses were infused at a constant rate over 30 minutes.

10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100

13.2 Animal Toxicology and/or Animal Pharmacology

Concretions consisting of the precipitated calcium salt of ceftriaxone have been found in the gallbladder bile of dogs and baboons treated with ceftriaxone. These appeared as a gritty sediment in dogs that received 100 mg/kg/day for 4 weeks.

A similar phenomenon has been observed in baboons but only after a protracted dosing period (6 months) at higher dose levels (335 mg/kg/day or more).

16 HOW SUPPLIED/STORAGE AND HANDLING

Ceftriaxone for Injection, USP is available in the following SmartPak® Pharmacy Bulk Package bags:

100 grams* One Pharmacy Bulk Package bag *Product No. 6100 NDC 66288-6100-1.

*Each 100 gram Pharmacy Bulk Package bag contains sterile ceftriaxone sodium equivalent to 100 grams of ceftriaxone

SmartPak® system components are not made with natural rubber latex.

Precautions

As with other cephalosporins, reconstituted Ceftriaxone for Injection tends to darken depending on storage conditions within the stated recommendations. However, product potency is not adversely affected. Use only if prepared solution is clear and free from particulate matter.

Storage Conditions

Prior to reconstitution, store Ceftriaxone for Injection sterile powder at room temperature 20°-25°C (68°-77°F) [see USP Controlled Room Temperature] and protected from light.

17 PATIENT COUNSELING INFORMATION

- Advise patients that neurological adverse reactions could occur with Ceftriaxone for Injection use. Instruct patients or their caregivers to inform their healthcare provider at once of any neurological signs and symptoms, including encephalopathy (disturbance of consciousness including somnolence, lethargy, and confusion), seizures, myoclonus, and non-convulsive status epilepticus, for immediate treatment, or discontinuation of Ceftriaxone for Injection [see *Warnings and Precautions* (5.3)].
- Patients should be advised that allergic reactions, including serious allergic reactions could occur and that serious reactions require immediate treatment and discontinuation of ceftriaxone. Patients should report to their health care provider any previous allergic reactions to ceftriaxone, cephalosporins, penicillins, or other similar antibacterials.
- Patients should be advised that diarrhea is a common problem caused by antibacterials, which usually ends when the antibacterial is discontinued. Sometimes after starting treatment with antibacterials, patients can develop watery and bloody stools (with or without stomach cramps and fever) even as late as two or more months after having taken the last dose of the antibacterial. If this occurs, patients should contact a physician as soon as possible.
- Patients should be counseled that antibacterial drugs including ceftriaxone, should only be used to treat bacterial infections. They do not treat viral infections (e.g., the common cold). When Ceftriaxone for Injection, USP is prescribed to treat a bacterial infection, patients should be told that although it is common to feel better early in the course of therapy, the medication should be taken exactly as directed. Skipping doses or not completing the full course of therapy may (1) decrease the effectiveness of the immediate treatment and (2) increase the likelihood that bacteria will develop resistance and will not be treatable by ceftriaxone or other antibacterial drugs in the future.

Rx only

SmartPak is a registered trademark of Samson Medical Technologies, L.L.C.

C6100e

Over a 0.15 to 3 g dose range in healthy adult subjects, the mean elimination half-life ranged from 5.8 to 8.7 hours, plasma clearance ranged from 0.58 to 1.45 L/hour, and renal clearance ranged from 0.32 to 0.73 L/hour.

Distribution

Ceftriaxone is reversibly bound to human plasma proteins and the binding of ceftriaxone decreases with increasing concentration from a value of 95% at plasma concentrations less than 25 mcg/mL to 85% at plasma concentration of 300 mcg/mL. Over a 0.15 to 3 g dose range in healthy adult subjects, the apparent volume of distribution ranged from 5.8 to 13.5 L.

Ceftriaxone crosses the blood placenta barrier.

Ceftriaxone penetrates the inflamed meninges of infants and pediatric patients. The average values of maximum plasma concentration, cerebrospinal fluid (CSF) concentrations, elimination half-life, plasma clearance and volume of distribution after a 50 mg/kg intravenous dose and after a 75 mg/kg intravenous dose in pediatric patients suffering from bacterial meningitis are shown in Table 4.

	50 mg/kg I.V.	75 mg/kg I.V.
	Maximum Plasma Concentrations (mcg/mL)	216
Elimination Half-life (hr)	4.6	4.3
Plasma Clearance (mL/hr/kg)	49	60
Volume of Distribution (mL/kg)	338	373
CSF Concentration – inflamed meninges (mcg/mL)	5.6	6.4
Range (mcg/mL)	1.3-18.5	1.3-44
Time after dose (hr)	3.7 (± 1.6)	3.3 (± 1.4)

After a 1 gram intravenous dose, average concentrations of ceftriaxone, determined from 1 to 3 hours after dosing, were 581 mcg/mL in the gallbladder bile, 788 mcg/mL in the common duct bile, 898 mcg/mL in the cystic duct bile, 78.2 mcg/gram in the gallbladder wall compared to a corresponding concentration of 62.1 mcg/mL in plasma.

Excretion

Ceftriaxone concentrations in urine are shown in Table 5.

Dose/Route	Average Urinary Concentrations (mcg/mL)					
	0-2 hr	2-4 hr	4-8 hr	8-12 hr	12-24 hr	24-48 hr
0.5 gram I.V.	526	366	142	87	70	15
1 gram I.V.	995	855	293	147	132	32
2 grams I.V.	2692	1976	757	274	198	40

Thirty-three percent to 67% of a ceftriaxone dose was excreted in the urine as unchanged drug, and the remainder was secreted in the bile and ultimately found in the feces as microbiologically inactive compounds.

The elimination of ceftriaxone is not altered by probenecid.

Special Populations

Average pharmacokinetic parameters of ceftriaxone in healthy subjects, elderly subjects, subjects with renal impairment, and subjects with liver disease are summarized in Table 6. Compared to healthy adult subjects, the pharmacokinetics of ceftriaxone were only minimally altered in elderly subjects and in patients with renal or hepatic impairment; therefore, dosage adjustments are not necessary for these patients with ceftriaxone dosages up to 2 grams per day. Ceftriaxone was not removed to any significant extent from the plasma by hemodialysis. In 6 of 26 dialysis patients, the elimination rate of ceftriaxone was markedly reduced, suggesting that plasma concentrations of ceftriaxone should be monitored in these patients to determine if dosage adjustments are necessary [see *Dosage and Administration* (2.1) and *Warnings and Precautions* (5.7)].

Subject Group	Elimination Half-Life (hr)	Plasma Clearance (L/hr)	Volume of Distribution (L)
	Healthy Subjects*	5.8-8.7	0.58-1.45
Elderly Subjects (mean age, 70.5 yr)	8.9	0.83	10.7
Patients with Renal Impairment			
Hemodialysis Patients (0-5 mL/min)**	14.7	0.65	13.7
Severe (5-15 mL/min)	15.7	0.56	12.5
Moderate (16-30 mL/min)	11.4	0.72	11.8
Mild (31-60 mL/min)	12.4	0.70	13.3
Patients with Liver Disease	8.8	1.1	13.6

*Dose ranged from 0.15 to 3 grams

**Creatinine clearance.

Drug Interactions

Interaction with Calcium: Two *in vitro* studies, one using adult plasma and the other neonatal plasma from umbilical cord blood, have been carried out to assess interaction of ceftriaxone and calcium. Ceftriaxone concentrations up to 1 mM (in excess of concentrations achieved *in vivo* following administration of 2 grams ceftriaxone infused over 30 minutes) were used in combination with calcium concentrations up to 12 mM (48 mg/dL). Recovery of ceftriaxone from plasma was reduced with calcium concentrations of 6 mM (24 mg/dL) or higher in adult plasma or 4 mM (16 mg/dL) or higher in neonatal plasma. This may be reflective of ceftriaxone-calcium precipitation.

12.4 Microbiology

Mechanism of Action

Ceftriaxone is a bactericidal agent that acts by inhibition of bacterial cell wall synthesis. Ceftriaxone has activity in the presence of some beta-lactamases, both penicillinases and cephalosporinases, of Gram-negative and Gram-positive bacteria.

Mechanism of Resistance

Resistance to ceftriaxone is primarily through hydrolysis by beta-lactamase, alteration of penicillin-binding proteins (PBPs), and decreased permeability.

Interaction with Other Antimicrobials

In an *in vitro* study antagonistic effects have been observed with the combination of chloramphenicol and ceftriaxone.

Antibacterial Activity

Ceftriaxone has been shown to be active against most isolates of the following bacteria, both *in vitro* and in clinical infections described in the INDICATIONS AND USAGE section:

- Gram-negative bacteria**
 - Acinetobacter calcoaceticus*
 - Enterobacter aerogenes*
 - Enterobacter cloacae*
 - Escherichia coli*
 - Haemophilus influenzae*
 - Haemophilus parainfluenzae*
 - Klebsiella oxytoca*
 - Klebsiella pneumoniae*
 - Moraxella catarrhalis*
 - Morganella morganii*
 - Neisseria gonorrhoeae*
 - Neisseria meningitidis*
 - Proteus mirabilis*
 - Proteus vulgaris*
 - Pseudomonas aeruginosa*
 - Serratia marcescens*

CEFTRIAXONE FOR INJECTION, USP

Manufactured for	
	Cherry Hill, NJ 08003, USA
	by
	ACS Dobfar S.p.A.
	20067 Tribiano (Milano) Italy
Printed in USA	
	6628861001